Draft version October 16, 2012
Preprint typeset using LaTeX style emulateapj v. 5/2/11

arXiv:1210.4027v1 [astro-ph.CO] 15 Oct 2012

THE HOST GALAXY OF THE SUPER-LUMINOUS SN 2010GX AND LIMITS ON EXPLOSIVE ^{56}NI PRODUCTION

Ting-Wan Chen[1]; Stephen J. Smartt[1]; Fabio Bresolin [2]; Andrea Pastorello[3]; Rolf-Peter Kudritzki [2]; Rubina Kotak [1]; Matt McCrum [1]; Morgan Fraser [1]; Stefano Valenti [4,5]

Draft version October 16, 2012

ABSTRACT

Super-luminous supernovae have a tendency to occur in faint host galaxies which are likely to have low mass and low metallicity. While these extremely luminous explosions have been observed from $z = 0.1$ to 1.55, the closest explosions allow more detailed investigations of their host galaxies. We present a detailed analysis of the host galaxy of SN 2010gx ($z = 0.23$), one of the best studied super-luminous supernovae. The host is a dwarf galaxy ($M_g = -17.42 \pm 0.17$) with a high specific star formation rate. It has a remarkably low metallicity of $12 + \log(\mathrm{O/H}) = 7.5 \pm 0.1$ dex as determined from the detection of the [OIII] $\lambda 4363$ Å line. This is the first reliable metallicity determination of a super-luminous supernova host. We collected deep multi-epoch imaging with Gemini + GMOS between $200 - 550$ days after explosion to search for any sign of radioactive ^{56}Ni, which might provide further insights on the explosion mechanism and the progenitor's nature. We reach *griz* magnitudes of $m_{\mathrm{AB}} \sim 26$, but do not detect SN 2010gx at these epochs. The limit implies that any ^{56}Ni production was below that of SN 1998bw (a luminous type Ic SN that produced around $0.4\,M_{\odot}$ of ^{56}Ni). The low volumetric rates of these supernovae ($\sim 10^{-4}$ of the core-collapse population) could be qualitatively matched if the explosion mechanism requires a combination of low-metallicity (below $0.2\,Z_{\odot}$), high progenitor mass ($> 60\,M_{\odot}$) and high rotation rate (fastest 10% of rotators).

Subject headings: Supernovae: general — Supernovae: individual (SN 2010gx)

1. INTRODUCTION

Recently, the new generation of wide-field survey projects, such as the Panoramic Survey Telescope And Rapid Response System (Pan-STARRS-1) (Kaiser et al. 2010) and the Palomar Transient Factory (PTF) (Rau et al. 2009), have carried out searches for supernovae without an inherent galaxy bias. Quimby et al. (2011) identified a class of hydrogen-poor "super-luminous supernovae (SLSNe)" with typical $g-$band absolute magnitudes of around -21.5 , which is 10 to 100 times brighter than normal core-collapse SNe. Their intrinsic brightness has allowed them to be discovered in the Pan-STARRS1 survey between $z = 1 - 1.55$ (Chomiuk et al. 2011; Berger et al. 2012). Pastorello et al. (2010) studied one of the closest examples, SN 2010gx ($z = 0.23$), in detail showing it to transition to a normal type Ic SN at 40 days after peak, while multi-colour photometry rules out ^{56}Ni as powering the peak magnitude. Two physical mechanisms for SN 2010gx (and the supernovae like it from Quimby et al.; Chomiuk et al.) have been proposed. One is that the type Ic SN is boosted in luminosity by energy deposition from magnetar spin down (Kasen & Bildsten 2010). The second is the shock breakout through a dense wind (Chevalier & Irwin 2011), or shock-interaction of the SN ejecta with massive C-O shells (Blinnikov & Sorokina 2010).

Neill et al. (2011) showed that most SLSNe tend to occur in star-forming dwarf galaxies, which have high specific star formation rates. Their intrinsic rate is $10^{-8}\,\mathrm{Mpc}^{-3}\,\mathrm{year}^{-1}$; or 10^{-4} of the normal core-collapse SN rate (Quimby et al. 2011; Gal-Yam 2012). However beyond this we lack information on their explosion sites and what the progenitor star population may be. An important question is whether or not the apparent preference they have for dwarf host galaxies is related to requiring low metallicity progenitor stars. As low metallicity affects stellar structure and evolution through changes in stellar winds, opacity and rotation, quantitative measurements are required. In this letter we study the host galaxy of one of the nearest SLSNe, SN 2010gx, and search for evidence of late emission from radioactive ^{56}Ni decay.

2. OBSERVATIONS

We used Gemini South and North Observatories with Gemini Multi-Object Spectrograph (GMOS) to collect *griz* photometry between $200 - 550$ days after the maximum of SN 2010gx (Table 1). To check for the presence of residual supernova flux at $200 - 350$ days after peak, we aligned and rescaled all images to the same pixel grid using IRAF/GEOMAP and GEOTRAN packages and made the assumption that the final image taken in January 2012 only contains flux from the host galaxy. We performed imaging subtraction using the High Order Transform of PSF ANd Template Subtraction (HOTPANTS) software [6], to subtract the January 2012 images

---

[1] Astrophysics Research Centre, School of Mathematics and Physics, Queen's University Belfast, Belfast BT7 1NN, UK
[2] Institute for Astronomy, 2680 Woodlawn Drive, Honolulu, HI 96822, USA
[3] INAF-Osservatorio Astronomico di Padova, vicolo dell'Osservatorio 5, 35122 Padova, Italy
[4] Las Cumbres Observatory Global Telescope Network, Inc. Santa Barbara, CA 93117, USA
[5] Department of Physics, University of California Santa Barbara, Santa Barbara, CA 93106-9530, USA

[6] http://www.astro.washington.edu/users/becker/hotpants.html

from the earlier epochs. The host galaxy subtracts off almost perfectly and no SN signal is detected. To determine a limiting magnitude for the SN flux in the three epochs, we added 6 fake stars, one at the supernova position and five within a surrounding radius of 0.6 arcmin to each of the $200-350$ days images. We ran this procedure multiple times with a range of fake star magnitudes and repeated image subtraction with HOTPANTS. We determined 5σ and 3σ detection limits by requiring that we could visually detect a fake SN exactly at the position of SN 2010gx and that the measured standard deviation of the set of fake stars added was 0.2 and 0.3 magnitudes respectively. The resulting upper limits are in Table 1.

The apparent magnitudes (SDSS AB system) of the host galaxy, SDSS J112546.72-084942.0, are $g = 23.71 \pm 0.14$, $r = 22.98 \pm 0.10$, $i = 22.90 \pm 0.13$ and $z = 22.98 \pm 0.16$. These were obtained from aperture photometry (within IRAF/DAOPHOT) on the best seeing images, using a zero-point calibration achieved with 10 SDSS reference stars. As deep, pre-explosion images of the host galaxy were not available, we compared our photometry results to both SDSS catalog magnitudes, and our own photometry performed on the SDSS images. In the $g-$band, the SDSS catalog Petrosian magnitude for the object is 22.8 ± 0.5, while we measure 22.6 ± 0.6 mag in the SDSS image. The large uncertainty is due to the marginal detection. Our deep Gemini image shows the host resolved from a fainter neighbouring galaxy (which is unresolved from the host in SDSS). If we use a large aperture on our Gemini $g-$band image, to include flux from the neighbouring galaxy then we obtain $g = 23.29 \pm 0.16$ mag. Within the uncertainties, the SDSS and Gemini magnitudes are similar and there is no hint that the +550 days Gemini images are brighter than the SDSS images and contain SN flux.

Long slit spectroscopy of SN 2010gx was carried out on 5 June 2010 (Pastorello et al. 2010) with the Gemini South Telescope + GMOS. The B600 grating (ID $G5323$) was used (1800s exposure) to obtain a spectral coverage in the observer frame of $4300-6400$ Å (henceforth referred to as the “blue spectrum”). Although this spectrum contains flux from SN 2010gx, we fitted a high order polynomial (order 15) through the SN continuum and subtracted off this contribution leaving a flat spectrum for which emission lines between $3500-6000$ Å could be measured. We could not measure the Hα line in this spectrum as it was contaminated by a strong sky line and slit nodding was not employed. A second spectrum was obtained on 23 Dec. 2011 with GMOS at the Gemini North Telescope, using the R400 grating (ID $G5305$; 3×1800s exposures), to cover $5070-8500$ Å (henceforth referred to as the red spectrum). The use of an order-blocking filter ($GG455$) truncated the wavelength coverage in the blue. Detrending of the data, such as bias-subtraction, was established using standard techniques within IRAF. For the red spectra, individual exposures were nodded along the slit, and we used two-dimensional image subtraction to remove the sky background. The spectra were wavelength calibrated using daytime arcs (CuAr lamp) and flux calibrated with the spectrophotometric standard Feige 34.

For both red and blue spectra, a $1.5''$ slit was used, yielding a resolution (as measured from narrow night sky lines) of ~ 6 Å in the red, and ~ 4.5 Å in the blue. The combined blue and red spectra provide spectral coverage between 4300 and 8500 Å and the overlap region was used to ensure a uniform flux calibration. Three strong lines ([OIII] λ5007, [OIII] λ4959, and Hβ) lie within the overlap region. We measured the line fluxes in both spectra and applied a linear scaling to bring the blue spectrum into agreement with the red. The line fluxes of these three lines agreed to within 3% after this rescaling.

Line flux measurements were made after fitting a polynomial function to subtract the continuum, and Fig. 3 shows the final combined spectrum of the host galaxy with this continuum subtracted. We fitted Gaussian line profiles within a custom built IDL environment, fixing the full-width-half-maximum (FWHM) of the mean of the three strong lines; 6.03 Å and 4.45 Å for the red and blue spectra respectively (see Table 2). We further normalized the spectrum and determined the equivalent width of each feature. The uncertainty estimation is calculated using the expression from Pérez-Montero & Díaz (2003), derived from the equivalent width of the line and the rms of the continuum (also listed in Table 2).

3. ANALYSIS

3.1. ^{56}Ni mass estimation for SN 2010gx

The fact that SN 2010gx was not detected after the host galaxy image subtraction at epochs between +226 and +334 days allows us to determine an upper limit for the mass of ^{56}Ni ejected. This is somewhat uncertain as it relies on assuming that the gamma ray trapping in SN 2010gx behaves like in other Ic SNe in the nebular phase, and the late luminosity is directly proportional to the mass of ^{56}Co present at any epoch.

We measured SN 2010gx fluxes at the effective rest wavelength of the $ri-$ filters and used a luminosity distance of 1114.7 Mpc ($m_u = 40.24$). We determined the upper limit of the luminosities and compared with SN 1998bw(Clocchiatti et al. 2011), a Ic SN with a high ^{56}Ni mass, in Johnson V and Cousins R in the rest frame (see Fig. 2). The most interesting measurement is the $i-$band limit for SN 2010gx flux at +226 days, which is significantly below what we would expect if SN 1998bw were at this distance. If we take $0.4\,M_\odot$ as the representative ^{56}Ni mass for SN 1998bw (McKenzie & Schaefer 1999; Sollerman et al. 2000; Maeda et al. 2006), then the flux limit of SN 2010gx (0.8 that of SN 1998bw; Fig. 2), suggests an upper limit of about $0.3\,M_\odot$ of ^{56}Ni ejected by SN 2010gx.

3.2. Host galaxy metallicity and other properties

We carried out an abundance analysis of the host galaxy based on the determination of the electron temperature, and detection of the [OIII] λ4363 auroral line (see Bresolin et al. 2009 for further details). Firstly, the emission line fluxes were reddening corrected using the Seaton (1979) Milky Way law of $A_V = 3.17 \times E(B-V)$. We assumed case B recombination which requires an intrinsic line ratio of Hα/Hβ = 2.86 and Hγ/Hβ = 0.47 at $T_e = 10^4$ K (Osterbrock 1989). The observed ratio of Hα/Hβ = 3.88 yields an extinction coefficient c(Hβ)= 0.48, which we used to calculate reddening-corrected line fluxes. The observed line fluxes and errors are listed in Table 2.

We confidently detected the [OIII] $\lambda4363$ auroral line (S/N$\sim$ 6) in the GMOS spectrum (see Fig. 3), allowing us to estimate the electron temperature and calculate the oxygen abundance directly. As in Bresolin et al. (2009), we used a custom-written PYRAF script based on the IRAF/NEBULA package. The direct method of estimating oxygen abundances uses the ratio of the intensities of [OIII] $\lambda4959$, $\lambda5007$/[OIII] $\lambda4363$ lines to determine the electron temperature (T_e) in the ionized gas region. We find an oxygen abundance of $12+\log(O/H) = 7.46\pm0.10$ dex, remarkably low even for dwarf galaxies (see Fig. 3). Assuming a solar value of 8.69 ± 0.05 dex (Asplund et al. 2009), this implies that the host galaxy is 1.2 dex below solar abundance or just $0.06\,Z_\odot$.

The reddening-corrected, and $z-$corrected flux of Hα is 3.49×10^{-16} ($\mathrm{erg\,s^{-1}\,cm^{-2}}$), from which we determined the star formation rate (SFR) of the host to be $0.41\,M_\odot\,\mathrm{year}^{-1}$ from the calibration of Kennicutt (1998): SFR ($M_\odot\,\mathrm{year}^{-1}$)= $7.9\times\ 10^{-42}$ L(Hα) ($\mathrm{erg s\,s^{-1}}$). We also estimated the SFR from the [OII] $\lambda3727$ line flux, which gives $0.36 \pm 0.10\,M_\odot\,\mathrm{year}^{-1}$ (Kennicutt 1998) and is consistent with Hα measurement. Berger et al. (2012) used [OII] $\lambda3727$ to estimate the SFR of the ultra-luminous PS1-12bam host, which at $z = 1.566$ has Hα shifted to the NIR. The agreement between the SFR estimates from [OII] $\lambda3727$ and Hα for the host of SN 2010gx supports [OII] $\lambda3727$ being reliable for these types of galaxies at higher redshift. We detected the galaxy ISM absorption of the Mg II $\lambda\lambda2796/2803$ Å doublet in the Gemini spectrum of 1 Apr. 2010 (Pastorello et al. 2010). The equivalent widths are 1.6 ± 0.2 and 2.6 ± 0.3 Å respectively, giving a ratio $W_{2796}/W_{2803} = 0.6$. The strength of the line $\lambda2803$ component is comparable with that seen toward GRB lines of site and somewhat higher than observed for PS1-12bam (Berger et al. 2012). The fact that the line ratio is significantly smaller than 2 (the ratio of their oscillator strengths) suggests we are not on the linear regime, nor the square root regime on the curve of growth which prevents further use of the lines as physical diagnostics. Nevertheless the absorption line strengths are similar to PS1-12bam and GRB sightlines. The line centroids give identical redshifts ($z = 0.231$) to the [OIII] emission lines ($z = 0.230$) detected in the same spectrum, giving reassurance that the Mg II absorption does arise in the host.

The $g-$band absolute magnitude of the host galaxy is $M_g = -17.42\pm0.17$ (adopting a Hubble constant of $H_0 = 72\,\mathrm{km\,s^{-1}\,Mpc^{-1}}$, $\Omega_\mathrm{M} = 0.3$, $\Omega_\lambda = 0.7$, to obtain the luminosity distance 1114.7 Mpc; Wright 2006), applying the Milky Way extinction of 0.13 (Schlafly & Finkbeiner 2011)), a k-correction 0.27 ± 0.08 (Chilingarian et al. 2010) and a correction for internal dust extinction of ~ 0.5 mag. The latter comes from the measured intrinsic Hα/Hβ ratio and assuming $R_V = 3.16$ (applicable for an LMC environment ; Pei 1992)[7]

The observed photometric $g - r$, $r - i$ colour terms were corrected for Milky Way extinction (Schlafly & Finkbeiner 2011) and used to fit the SED of a stellar population model (Maraston 2005). We used the low-metallicity (Z$\sim$ $1/20\,Z_\odot$) models, assuming a Salpeter initial mass function with red horizontal branch morphology. The measured colours give reasonable agreement with a population model of age between 20 to 30 Myr, and a stellar mass around $6.17 \times 10^7\,M_\odot$. We also employed the MAGPHYS galaxy SED models of da Cunha et al. (2008). Fig. 1 shows the observed SED and the redshifted MAGPHYS best-fit model spectrum : total stellar mass of $1.58 \times 10^8\,M_\odot$ and hence a specific star formation rate (sSFR) of $2.59 \times 10^{-9}\,\mathrm{year}^{-1}$.

4. DISCUSSION AND CONCLUSION

It has already been demonstrated that the peak luminosity of super-luminous SNe such as SN 2010gx cannot be due to ^{56}Ni (Pastorello et al. 2010; Quimby et al. 2011; Chomiuk et al. 2011). Chomiuk et al. illustrated that applying Arnett's rule (Arnett 1982), results in an unphysical solution, with the mass of ^{56}Ni exceeding the total C+O ejecta mass. Our search for a luminous tail phase in SN 2010gx was not motivated by what powers the peak luminosity, rather it is to determine if there is any sign of radioactive ^{56}Co at this stage and any similarity to ^{56}Ni-rich Ic SNe such as SN 1998bw. The only robust conclusion we can draw is that there is unlikely to be any excessive mass of ^{56}Co above and beyond about $0.3\,M_\odot$. At a similar phase, SN 2010gx is fainter than SN 1998bw as shown in Fig. 2. This limit on the late luminosity will also allow restrictions on the magnetar models of Kasen & Bildsten (2010) when put in context with other super-luminous supernovae of similar types (Inserra et al. in prep).

Our detection of the [OIII] $\lambda4363$ auroral line gives confidence to our measurement of the remarkably low metallicity of $12+\log(O/H) = 7.46\pm0.10$ dex ($0.06\,Z_\odot$). In Fig. 3 we show a compilation of other dwarf galaxies with the [OIII] $\lambda4363$ direct method measurements, including four GRB hosts from Kewley et al. (2007). The host of SN 2010gx is on the lower end and is amongst the lowest metallicity dwarf galaxies known. It has been clear from the early discoveries of these SNe at $z < 1$ (Pastorello et al. 2010; Quimby et al. 2011; Chomiuk et al. 2011), that the hosts are faint dwarf galaxies and Neill et al. (2011) showed that they have high sSFRs. If the extremely low value we determine is a common factor amongst these SNe then it will be a major constraint on the progenitor channel.

The magnetar scenario suggested by Kasen & Bildsten (2010) to explain these SNe may benefit from low metallicity progenitors. Massive stars rotate more rapidly at SMC metallicity ($0.2\,Z_\odot$) than solar (Hunter et al. 2008 ; Martayan et al. 2007), although $0.06\,Z_\odot$ is unexplored territory. Evolutionary models of rotating stars have focused attention on low metallicity environments as mass-loss is reduced (Mokiem et al. 2007) hence angular momentum loss is lower (Maeder & Meynet 2000). The energy input required from a magnetar requires the neutron star to be both highly magnetic ($B \sim 10^{14}$ G) and rotating rapidly at formation ($P_i \sim 2 - 20$ ms). It must also occur in a Wolf-Rayet or carbon-oxygen star, since there is no sign of hydrogen in the spectra of SN 2010gx or any similar SNe. Gal-Yam (2012) gives an estimated relative rate of the number of 2010gx-like SNe compared to the total core-collapse SN rate of 10^{-4}. If the progenitor channel requires low metallicity (as we measure), high mass (to produce a WC star) and fast rotation (to produce a rapidly rotating magnetar) then this very low rate may not be unexpected. Young et al. (2008) estimate that ~1% of the star formation in the Local Universe ($z < 0.04$) is in galaxies with $Z < 0.2\,Z_\odot$; around 13% of massive stars above the SN threshold of $8\,M_\odot$ are $> 60\,M_\odot$ (assuming a Salpeter IMF); and about 10% of O-stars in the SMC rotate faster than $\sim$300 km s^{-1} (Mokiem et al. 2007). One can speculate that all of these three conditions are required to produce the rate of the 2010gx-like SNe and the estimated rate (which is itself uncertain) would be qualitatively reproduced. The high-mass requirement could also be substituted for an interacting binary system which causes spin-up in the CO-core that collapses, similar to the proposed GRB systems (Cantiello et al. 2007). The Chevalier & Irwin (2011) scenario of interaction with a dense wind would probably require extreme progenitor mass and pulsational mass-loss. The fact that we see Wolf-Rayet stars at very low metallicity (e.g. in IZw13 at $0.02\,Z_\odot$) (Brown et al. 2002) could suggest that pulsational mass-loss is more prevalent at low metallicity. This one measurement is intriguing new information on these extreme SNe but further measurements of the metallicity, star formation rate and volumetric rates of these supernovae are required to constrain their explosion channels further.

---

[7] This would be reduced by 0.05 mag if we used a value suitable for an SMC environment; $R_V = 2.93$ (Pei 1992).

ACKNOWLEDGMENTS

S.J.S acknowledges support from the European Research Council in the form of an Advanced Grant. T.-W. Chen appreciates to Meng-Chung Tsai, Zheng Zheng and Kai-Lung Sun for valuable comments and Prof. Christy Tremonti, Dr. Elisabete da Cunha and Dr. Stephane Charlot for useful advice; thanks to all QUB SN group members for discussions.

Facilities: Gemini North and South (GMOS)

REFERENCES

Amorín, R., Pérez-Montero, E., Vílchez, J. M., & Papaderos, P. 2012, ApJ, 749, 185
Arnett, W. D. 1982, ApJ, 253, 785
Asplund, M., Grevesse, N., Sauval, A. J., & Scott, P. 2009, ARA&A, 47, 481
Berg, D. A., Skillman, E. D., Marble, A. R., et al. 2012, ApJ, 754, 98
Berger, E., Chornock, R., Lunnan, R., et al. 2012, ApJ, 755, L29
Blinnikov, S. I., & Sorokina, E. I. 2010, ArXiv e-prints
Bresolin, F., Gieren, W., Kudritzki, R.-P., et al. 2009, ApJ, 700, 309
Brown, T. M., Heap, S. R., Hubeny, I., Lanz, T., & Lindler, D. 2002, ApJ, 579, L75
Cantiello, M., Yoon, S.-C., Langer, N., & Livio, M. 2007, A&A, 465, L29
Chevalier, R. A., & Irwin, C. M. 2011, ApJ, 729, L6
Chilingarian, I. V., Melchior, A.-L., & Zolotukhin, I. Y. 2010, MNRAS, 405, 1409
Chomiuk, L., Chornock, R., Soderberg, A. M., et al. 2011, ApJ, 743, 114
Clocchiatti, A., Suntzeff, N. B., Covarrubias, R., & Candia, P. 2011, AJ, 141, 163
da Cunha, E., Charlot, S., & Elbaz, D. 2008, MNRAS, 388, 1595
Gal-Yam, A. 2012, Science, 337, 927
Gal-Yam, A., Mazzali, P., Ofek, E. O., et al. 2009, Nature, 462, 624
Guseva, N. G., Papaderos, P., Meyer, H. T., Izotov, Y. I., & Fricke, K. J. 2009, A&A, 505, 63
Hoyos, C., Koo, D. C., Phillips, A. C., Willmer, C. N. A., & Guhathakurta, P. 2005, ApJ, 635, L21
Kaiser, N., Burgett, W., Chambers, K., et al. 2010, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 7733, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series
Kasen, D., & Bildsten, L. 2010, ApJ, 717, 245
Kennicutt, Jr., R. C. 1998, ARA&A, 36, 189
Kennicutt, Jr., R. C., Bresolin, F., & Garnett, D. R. 2003, ApJ, 591, 801
Kewley, L. J., Brown, W. R., Geller, M. J., Kenyon, S. J., & Kurtz, M. J. 2007, AJ, 133, 882
Kudritzki, R.-P., Urbaneja, M. A., Gazak, Z., et al. 2012, ApJ, 747, 15
Lee, H., Skillman, E. D., Cannon, J. M., et al. 2006, ApJ, 647, 970
Lee, J. C., Salzer, J. J., & Melbourne, J. 2004, ApJ, 616, 752
Maeda, K., Nomoto, K., Mazzali, P. A., & Deng, J. 2006, ApJ, 640, 854
Maeder, A., & Meynet, G. 2000, ARA&A, 38, 143
Maraston, C. 2005, MNRAS, 362, 799
Martayan, C., Frémat, Y., Hubert, A.-M., et al. 2007, A&A, 462, 683
McKenzie, E. H., & Schaefer, B. E. 1999, PASP, 111, 964
Modjaz, M., Kewley, L., Kirshner, R. P., et al. 2008, AJ, 135, 1136
Mokiem, M. R., de Koter, A., Vink, J. S., et al. 2007, A&A, 473, 603
Neill, J. D., Sullivan, M., Gal-Yam, A., et al. 2011, ApJ, 727, 15
Osterbrock, D. E. 1989, Astrophysics of gaseous nebulae and active galactic nuclei
Pagel, B. E. J., Edmunds, M. G., Blackwell, D. E., Chun, M. S., & Smith, G. 1979, MNRAS, 189, 95
Papaderos, P., Guseva, N. G., Izotov, Y. I., & Fricke, K. J. 2008, A&A, 491, 113
Pastorello, A., Smartt, S. J., Botticella, M. T., et al. 2010, ApJ, 724, L16
Pei, Y. C. 1992, ApJ, 395, 130
Pérez-Montero, E., & Díaz, A. I. 2003, MNRAS, 346, 105
Quimby, R. M., Kulkarni, S. R., Kasliwal, M. M., et al. 2011, Nature, 474, 487
Rau, A., Kulkarni, S. R., Law, N. M., et al. 2009, PASP, 121, 1334
Richer, M. G., & McCall, M. L. 1995, ApJ, 445, 642
Schlafly, E. F., & Finkbeiner, D. P. 2011, ApJ, 737, 103
Seaton, M. J. 1979, MNRAS, 187, 73P
Skillman, E. D., Kennicutt, R. C., & Hodge, P. W. 1989, ApJ, 347, 875
Sollerman, J., Kozma, C., Fransson, C., et al. 2000, ApJ, 537, L127
Wright, E. L. 2006, PASP, 118, 1711
Young, D. R., Smartt, S. J., Mattila, S., et al. 2008, A&A, 489, 359
Young, D. R., Smartt, S. J., Valenti, S., et al. 2010, A&A, 512, A70



Fig. 1.— *Left* : upper- Colour combined (*gri*) Gemini GMOS images of the host galaxy of SN 2010gx; middle- A fake SN 5σ detection in r-band after mage subtraction; bottom- A fake 3σ detection. *Right*: Photometry of the host galaxy (red circle) with the best-fit ($\chi^2 = 0.58$) model galaxy SED (black line) from MAGPHYS (da Cunha et al. 2008), the blue line shows the same model without attenuation by dust.



FIG. 2.— The limiting luminosities of SN 2010gx compared with SN 1998bw (Clocchiatti et al. 2011). At 226 days we have a flux limit of 0.8 that of SN 1998bw.



FIG. 3.— *Upper*: Spectrum of the host galaxy of SN 2010gx. The [OIII] λ4363 Å auroral line is shown in the inset. *Bottom*: Luminosity-metallicity relationship for dwarf galaxies in the local Universe, showing the metallicity as measured using the direct (Te) method, and using data from the literature listed in the figure key. Grey lines are the boundary of normal galaxy mass-metallicity relation adapted from Kudritzki et al. (2012), where metallicities have been calculated using blue supergiants.

Date	MJD	Phase	g	r	i	z	Telescope	Instrument
2010 Jun 4	55351.97	56.5			21.13(0.19)		GS	GMOS-S
2010 Jun 8	55354.93	58.9	23.55(0.28)	21.52(0.09)	21.31(0.12)		WHT	ACAM
2010 Jun 8	55355.48	59.3		21.58(0.13)	21.31(0.16)		FTS	EM03
2010 Jun 13	55360.47	63.4	23.91(0.30)	21.81(0.20)	21.56(0.13)		FTS	EM03
2010 Jun 13	55360.92	63.8	23.95(0.49)	21.83(0.12)	21.58(0.29)		LT	RATcam
2010 Jun 16	55363.48	65.8	>23.73	21.91(0.16)			FTS	EM03
2010 Jun 29	55376.38	76.3		22.47(0.21)	22.38(0.33)		FTS	EM03
2010 Dec 30	55560.63	226.1			H 22.90(0.13) >25.94(0.17)	H 22.98(0.16) >24.58(0.29)	GN	GMOS-N
2011 Jan 14	55575.32	238.1	H 23.71(0.14) >26.42(0.18)	H 22.98(0.10) >25.68(0.11)			GS	GMOS-S
2011 Feb 28	55620.12	274.5			H 23.20(0.13) >24.78(0.13)	H 23.00(0.13) >23.35(0.32)	GS	GMOS-S
2011 May 20	55701.28	340.5	H 23.72(0.13) >26.00(0.15)				GN	GMOS-N
2011 May 25	55706.08	344.4		H 23.08(0.08) >25.35(0.15)	H 23.02(0.11) >24.60(0.20)	H 23.06(0.16)	GS	GMOS-S
2012 Jan 18	55944.22	538.0		H 23.03(0.10)			GS	GMOS-S
2012 Jan 20	55946.23	539.6			H 23.17(0.11)		GS	GMOS-S
2012 Jan 21	55947.21	540.4	H 23.62(0.11)				GS	GMOS-S

TABLE 1

Observed photometry of SN 2010gx and its host galaxy. Data until 2010 Dec 30 are from Pastorello et al. (2010). The symbol "H" indicates host galaxy photometry, and the ">" expresses the limiting magnitude of the SN (5σ). Phase has been corrected for time dilation, based on the SN explosion at MJD 55282.51.

SDSS name	SDSS J112546.72-084942.0
RA	11:25:46.72
Dec	-08:49:42.0
Redshift	0.23
Apparent g (mag)	23.71 ± 0.14
Apparent r (mag)	22.98 ± 0.10
Apparent i (mag)	22.90 ± 0.13
Apparent z (mag)	22.98 ± 0.16
Galactic extinction$_g$ (mag)	0.13
k-correction$_g$ (mag)	0.27 ± 0.08
Internal extinction$_g$ (mag)	$\sim$ 0.5 (R_V = 3.16)
Luminosity distance (Mpc)	1114.7
Absolute g (mag)	-17.42 ± 0.17
Physical Diameter (kpc)	3.1
Hα Flux (erg s^{-1} cm^{-2})	3.49e-16
SFR ($M_\odot$ year^{-1})	0.41
Stellar mass ($M_\odot$)	$10^{8.17}$
sSFR (year^{-1})	2.76×10^{-9}
$12 + \log(\mathrm{O/H})$	7.46 ± 0.1
Line	Measured Flux ± Error (erg s^{-1} cm^{-2})
[OII] λ3727	1.297e-16 ± 4.33e-18
[NeIII] λ3868	6.120e-17 ± 2.65e-18
Hδ λ4102	4.254e-17 ± 2.97e-18
Hγ λ4340	5.267e-17 ± 3.59e-18
[OIII] λ4363	2.269e-17 ± 2.33e-18
Hβ λ4861	1.285e-16 ± 3.25e-18
[OIII] λ4959	2.129e-16 ± 3.37e-18
[OIII] λ5007	6.569e-16 ± 8.13e-18
Hα λ6563	4.996e-16 ± 5.07e-18
[NII] λ6584	9.807e-18 ± 8.21e-19
[SII] λ6717	1.918e-17 ± 7.96e-19
[SII] λ6731	1.820e-17 ± 9.29e-19
c(Hβ)	0.48 ±0.04
-W(Hβ) ($\AA$)	52

TABLE 2
Main properties of the host galaxy of SN 2010gx.